CALENDAR

OF

FLORA,

COMPOSED DURING THE YEAR

1809,

at

Marrington,

Lat. 53°. 30'.

BY GEORGE CROSFIELD, SECRETARY

TO THE BOTANICAL SOCIETY OF WARRINGTON.

These, as they change, ALMIGHTY FATHER, these, Are but the varied God. The rolling year Is full of Thee.------

Mysterious round! what skill, what force divine, Deep felt, in these appear! a simple train, Yet so delightful mix'd, with such kind art, Such beauty and beneficence combin'd; Shade, unperceiv'd, so softening into shade; And all so forming an harmonious whole; That, as they still succeed, they ravish still.

THOMSON.

WARRINGTON:
Printed by J. Haddock.

SOLD BY WILKIE AND ROBINSON, PATERNOSTER-ROW;
AND WHITE, FLEET-STREET,
LONDON.

PREFACE.

SINCE the science of Botany became a branch of study, considered at once useful and instructive, it has been deemed of some importance to point out the different vegetable productions of particular districts. Some very 'respectable Botanists have likewise expressed a wish that the periods of infloreseence in different seasons and in different latitudes should be marked down with accuracy. To unite these two plans, at the least possible expence, is my design in the following pages, which contain a list of upwards of eight hundred British Plants, eomprized in the twenty-three first Classes of the Linnean system, growing, either wild or in a cultivated state, in the neighbourhood of Warrington, and arranged according to their earliest periods of infloreseence during the year 1809. It would, I am aware, have added much to its value, to have been accompanied by an accurate meteorological journal; but such an undertaking would have oeeupied more time than was consistent with my commercial avocations. It may not however be amiss to say, that, with the exception of a few warm days early in the 3d Month, (March.) the Spring was delayed later than usual by constant easterly winds, vegetation being kept back until the third of the 5th Month, (May.). From this time until the beginning of the following month the progress of vegetation was very rapid. The latter end of Summer was attended with much rain.

In compiling the following Calendar of Flora, I have not only depended on my own personal observation, but have availed myself of the advantages afforded by my situation as

Secretary to a provincial Botanical Society, to some of the members of which I aeknowledge myself under considerable obligation for their ready assistance.

With a few exceptions, no plant has been admitted into the following list, which is not ranked as indigenous in the very excellent *Flora Britannica* of Dr. Smith. Where the rule is departed from, the authority for it is specified.

An asterisk has been prefixed to all those plants, which, although indigenous, are only found, in this neighbourhood, in a state of cultivation. A few, observed in the more western, and northern parts of this county, are distinguished by a dagger.

The notes will not I trust, be deemed superfluous. For those marked K, I am indebted to my esteemed friend James Kendrick, M. D. F. L. S. President of the Botanical Society of Warrington.

The town of Warrington is situated at the southern extremity of the county of Laneaster, the river Mersey dividing it from that of Chester. The broad estuary of this River commences about six miles westward. In its immediate vicinity, is a large quantity of rich meadow land, formed principally of alluvial soil. A large extent of peat moss commences about two miles eastward of the town. The only high ground in its neighbourhood is a hill about two miles to the south, formed of red sand stone, covered with a stratum of sandy soil.

In every instance I have adopted the nomenclature of Dr. Smith: where he has differed from Linnæus, or where subsequent writers have differed from him their synonyms are also given.

GEORGE CROSFIELD.

WARRINGTON,
1st of 1st Month, 1810.

A CALENDAR, &c.

FIRST MONTH (January).

- 14. ULEX europæus. Common Furze.
- 16. BELLIS perennis. Common Daisy. (1)

SECOND MONTH (February).

- 2. * HELLEBORUS fatidus. Stinking Hellebore.
- 12. SENECIO vulgaris. Groundsel.
- 14. GALANTHUS nivalis. Snowdrop.

 * CROCUS vernus. Spring Crocus.

 VINCA minor. Lesser Periwinkle.
- 26. DAPHNE Mezereum. Mezereon.

THIRD MONTH (March).

- 2. LAMIUM purpureum. Red Archangel.
- 7. \$\beta \text{VIOLA odorata.} \text{Sweet Violet.}
- 10. PRIMULA vulgaris.

 y PRIMULA veris. Linn. Primrose.

- 11. LEONTODON Taraxacum. Common Dandelion. (2)
 STELLARIA media.
 ALSINE media. Linn. Common Chickweed.
- 12. * SCILLA bifolia. Two-leaved Squill.

 RANUNCULUS Ficaria. Pilewort. (3)

 BETULA Alnus.
 ALNUS glutinosa. Willdenow. Common Alder.

 CORYLUS Avellana. Hasel-nut-tree..

 POPULUS alba. Great White Poplar.
- 13. ULMUS campestris. ULMUS suberosa. Willd. Common Elm-tree.
- 14. FRAGARIA sterilis. Barren Strawberry. SALICIS species varia. Various kinds of Willow. (4)
- POA annua. Annual Meadow-grass. DRABA verna. Common Whitlow-grass.
- 19. * PULMONARIA officinalis. Common Lung-wort.
- a VIOLA odorata. Sweet Violet.
 NARCISSUS Pseudo-narcissus. Common Daffodil.
 DAPHNE Laureola. Spurge-Laurel.
 CARDAMINE hirsuta. Hairy Ladies-Smock.
- 25. VERONICA verna. Vernal Speedwell. OXALIS Acetosella. Common Wood-Sorrel.
- (2) Whether this plant possesses the deobstruent properties attributed to it by some authors of reputation may be doubted. Its diuretic effects are in some instances considerable. Its vulgar name, in more than one language, implies a knowledge of its powers. K_{\bullet}
- (3) It may not be amiss to state that this termination so common in the names of British plants, is derived from the Saxon Wyrt, a herb or plant, and implies that the plant so denominated, has been supposed to possess some specific power of curing the disease of a particular organ, e, g. Lungwort, Liverwort, &c, &c, K.
- (4) The bark of all the species of Salix, containing a considerable portion of Tannin, is useful in the art of tanning leather. With regard to its medical properties, it may suffice to say, that it is by no means a despicable tonic, though experience and analysis prove it an improper substitute for the Bark of the Cinchona lancifolia (Peruvian Burk). K.

25. ULMUS montana. Willd. Broad-leaved Elm-tree.

26. β VIOLA tricolor. Pansy Violet.

ADOXA Moschatellina. Tuberous Moschatel.

ANEMONE nemorosa. Wood Anemone. has a first THLASPI arvense. Smooth Mithridate Mustard.

MERCURIALIS perennis. Perennial Mercury.

27. * SCILLA verna. Vernal Squill.

JUNCUS pilosus. Hairy Rush.

RIBES Grossularia. Rough Gooseberry.

LAMIUM album. White Archangel.

COCHLEARIA officinalis. Common Scurvy-grass.

POPULUS tremula. Aspen.

nigra. Black Poplar.

TAXUS baccata. Yew Tree.

28. MONTIA fontana. Water Chickweed.

PRIMULA elatior.
β PRIMULA veris. Linn. Oxlip.

CHRYSOPLENIUM oppositifolium. Opposite-leaved
Golden-Saxifrage.

RANUNCULUS hederaccus. Ivy Crowfoot.

CALTHA palustris. Marsh Marygold.

FOURTH MONTH (April).

CHEIRANTHUS fruticulosus. Wallflower.

1. VERONICA hederifolia. Ivy-leaved Speedwell.

ALCHEMILLA arvensis.
APHANES arvensis. Linn.
ALCHEMILLA aphanes. Willd.

CERASTIUM vulgatum. Broad-leaved Mouse-car
Chickweed.
GLECHOMA hederacea. Ground-Ivy.
ARABIS thaliana. Common Wall-Cress.

2. VIOLA cauina. Dog's Violet.

JUNCUS sylvaticus.

JUNCUS pilosus. Linn.

Wood Rush.

3. CHARA vulgaris. Common Chara. * TULIPA sylvestris. Wild Tulip.

- *ORNITHOGALUM luteum. YellowStar of Bethlehem. 3.
 - * ANEMONE apennina. Blue mountain Anemone.
 - * FUMARIA solida. Solid bulbous Fumitory.

THLASPI campestre, Common Mithridate Mustard.

BUXUS sempervirens. Box-tree.

- * RUSCUS aculeatus. Butcher's-Broom.
- * VISCUM album. Misseltoe.
- 4. * VINCA major. Greater Periwinkle.
- 5. RIBES rubrum. Common Currants.
 - * FRITILLARIA Meleagris. Common Fritillary.
 - * ANEMONE Pulsatilla. Pasque Flower. MYRICA Gale. Sweet Gale.
- PRIMULA farinosa. Bird's eye Primrose. 7.
- * HELLEBORUS viridis. Green Hellebore. 8.
- GALIUM Aparine. Goose-grass. (5) 9.
- RIBES Uva-crispa. Smooth Gooseberry. 10. CARPINUS Betulus. Horn-beam. POPULUS canescens. Common White Poplar. * ECHIUM vulgare. Common blue Vipers-Bugloss.
- 11. TUSSILAGO hybrida. Longer stalked Butter-bur. * ARABIS stricta. Bristol Rock-cress.
- α CALLITRICHE aquatica. Water Star-wert. CALLITRICHE verna. Linn. 14.

ALCHEMILLA vulgaris. Common Ladies Mantle. * DENTARIA bulbifera. Bulbiferous Coral-wort.

- CARDAMINE pratensis. Meadow Ladies-smock.
- PRIMULA veris. Common Covislip. 15. α VIOLA tricolor. Pansy Violet. PRUNUS spinosa. Sloe-tree. As 1 41.
- ----- insititia. Bullace-tree. 17.
 - * SCROPHULARIA vernalis. Yellow Fig-wort.

⁽⁵⁾ The powers of late attributed to this plant, as an anticarcinomatous remedy, may reasonably be doubted. It has long been a popular remedy in those complaints commonly called scorbutic. Its root, in common with that of Madder, (Rubia tinctorum Linn.) has the property of tinging the bones of animals fed upon it, of a red colour. K.

- 17. * COCHLEARIA danica. Danish Scurvy-grass. ERYSIMUM Alliaria. Garlick Hedge-mustard. Re-
- * NARCISSUS poeticus. Poetic Narcissus. 18. PRUNUS domestica. Plum-tree.
- 19. CHARA hispida. Prickly Chara. Red on RANUNCULUS auricomus. Wood Crowfoot.
- CERASTIUM aquaticum. Water Chickweed. PRUNUS Cerasus. Cherry-tree. THLASPI Bursa-Pastoris. Common Shepherd's Purse. BRASSICA Rapa. Turnep. 7 . . .
- * IXIA Bulbocodium. Crocus-leaved Ixia. (Curtis.) 25.
 - * EPIMEDIUM alpinum. Barren-wort.
 - * NARCISSUS biflorus. Pale Narcissus. PYRUS communis. Iron Pear-tree.
- RIBES nigrum. Black Currants. 26. * ASARUM europæum. Asarabacca.
- 27. CHÆROPHYLLUM sylvestre. Smooth Cow Parsley. STELLARIA nemorum. Wood Stitchwort. COCHLEARIA anglica. English Scurvy-grass. SALIX repens. Creeping dwarf Willow.
- STELLARIA holostea. Greater Stitchwort. 28. BETULA alba. Common Birch. 774
- VERONICA arrensis. Wall Speedwell. 29. ASPERULA odorata. Sweet Woodruff. EUPHORBIA Cyparissias. Cypress Spurge. Level of FRAGARIA vesca. Wood Strawberry.
 - a RANUNCULUS aquatilis. Water Crowfoot β RANUNCULUS aquatilis.
- α VALERIANA Locusta.
 VALERIANA olitoria. Willd. } Corn Sallad. \$0. ERIOPHORUM vaginatum. Single-headed Cotton-grass. ----- angustifolium. Common Cotton-grass. PLANTAGO lanceolata. Ribwort Plantain. (6) President

⁽⁶⁾ A year or two ago, I met with a variety of this species, having the summit of the spike surmounted by leaves; which I considered as an accidental conversion of the terminal flowers, rather than as a permanent variety.

Qu. Is this var. 3. of Withering, described as having "the summit of the stalk surrounded with leaves longer than the spike?"

* ERICA Dabeoci.

** ERICA Dabeoci.

MENZIESIA polifolia. Jussieu.

ANDROMEDA polifolia. Marsh Rosemary.

** LYCHNIS dioica. Red Campion.

IBERIS nudicaulis. Naked-stalked Candy-tuft.

TURRITIS hinsma. Hairy Tower-Mustard.

SINAPIS arvensis. Charlock.

GERANIUM molle. Dove's-foot Crane's-bill.

ORCHIS mascula. Early purple Orchis.

FIFTH MONTH (May).

- 1. VERONICA agrestis. Procumbent Speedwell.

 ANCHUSA sempervirens. Evergreen Alkanet.

 ACER Pseudo-platanus. Greater Maple.

 CERASTIUM semidecandrum. Little Mouse-ear-Chickweed.
 - * TROLLIUS europaus. Globe-flower.
 - * FUMARIA lutea. Yellow Fumitory.
 - * SALIX herbacea. Least Willow.
- 2. * GENTIANA acaulis. Dwarf Gentian. (E. Bot.)
- 3. VACCINIUM Myrtillus. Bilberry.
 ACER campestre. Common Maple.
 HOLOSTEUM umbellatum. Umbelliferous Chickweed.
 CARDAMINE amara. Bitter Ladies-smock.
 ERODIUM cicutarium.
 GERANIUM cicutarium. Linn. Ilemlock Stork's-bill.
 GERANIUM pusillum. Small-flowered Crane's-bill.
 FAGUS sylvatica. Beech-tree.
- 5. HIPPURIS vulgaris. Common Mare's-tail. (7)
 - * ORNITHOGALUM umbellatum. Common Star of Bethlehem.

GERANIUM lucidum. Shining Crane's-bill. CAREX pracox. Vernal Carex.

⁽⁷⁾ Was observed to emerge on the 12th of 4th Month. (April.) K.

- 6. * ARBUTUS Uva-ursi. Red-berried trailing Arbutus. (8)
 - * SAXIFRAGA umbrosa. London Pride.
- 7. BUNIUM flexuosum. Common Earth-nut.
 - * SAXIFRAGA oppositifolia. Purple Saxifrage.

MESPILUS Oxyacantha. CRATÆGUS Oxyacantha. Linn. } Hawthorn. (9)

CAREX paludosa. Lesser common Carex.

8. VERONICA serpyllifolia. Smooth Speedwell.

* VIOLA lutea. Yellow mountain Pansy.

SCANDIX odorata. Great Chervil.

PYRUS Malus. Apple-tree. 1. 4

* GEUM rivale. Water Avens.

PEDICULARIS palustris. Marsh Louse-wort.

----- sylvatica. Pasture Louse-wort. (10)

POLYGALA vulgaris. Milk-wort.

CAREX dioica. Common separate-headed Carex.

* SALIX reticulata. Wrinkled Willow.

9. VERONICA montana. Mountain Speedwell.

ANTHOXANTHUM odoratum. Sweet-scented Vernal-grass.

SAGINA apetala. Annual small-flowered Pearl-wort.

CHENOPODIUM Bonus Henricus. Perennial Goosefoot.

SCILLA nutans.
HYASCINTHUS nonscriptus. Linn. Hare-bell.

* SAXIFRAGA hypnoides. Mossy Saxifrage.

* CERASTIUM tomentosum. Woolly Mouse-ear. (Hull.)

ANTIRRHINUM Cymbalaria. Ivy-leaved Snapdragon.

⁽⁸⁾ The tonic and sedative properties of this plant are not, I apprehend, in general fully appreciated, notwithstanding what has been said on the subject by Doctors Ferriar and Bourne. In many cases I have found it not a little efficacious. K.

⁽⁹⁾ Its inflorescence usually commences on the first of this month (MAY).

⁽¹⁰⁾ Frequently varies to white,

9.	* IBERIS amara. Bitter Candy-tuft.
υ.	FUMARIA officinalis. Common Fumitory.
	* GNAPHALIUM dioicum. Mountain Cud-weed.
10.	
10.	RANUNCULUS sceleratus. Celery-leaved Crowfoot.
	hirsutus. Pale hairy Crowfoot.
	acris. Upright meadow Crowfoot.
*	GALEOBDOLON luteum. GALEOPSIS Galeobdolon. Linn. Yellow Dead-Nettle.
	LEONURUS Galcobdolon. Willd.)
	* GERANIUM phaum. Dusky Crane's-bill.
	* CENTAUREA Cyanus. Com Blue-bottle.
-11.	FRAXINUS excelsior. Common Ash.
	VERONICA Chamedrys. Germander Speedwell.
	VALERIANA dioica. Marsh Valerian.
	SHERARDIA arvensis. Blue Sherardia.
	β MYOSOTIS scorpioides. Monse-ear Scorpion-grass.
	MENYANTHES trifoliata. Buckbean. (11)
	* LONICERA Caprifolium. Pale perfoliate Honeysuckle.
	GLAUX maritima. Black Salt-wort.
	STATICE Armeria. Common Thrift.
	JUNCUS campestris. Hairy field Rush.
	RUMEX Acetosa. Common Sorrel.
	TRIGLOCHIN maritimum. Sea Arrow-grass
	FARIS quantous. There I mass
	ARENARIA rubra. Purple Sandwort.
	PRUNUS Padus. Bird Cherry.
	PYRUS aucuparia. Linn Mountain-ash.
	SORBUS aucuparia. Linn. Mountain-ash.

---- reptans. Trailing Tormentil.

Comman Tormentil.

POTENTILLA anserina. Silver-weed.

TORMENTILLA erecta. Linn.

TORMENTILLA officinalis.

⁽¹¹⁾ Varies frequently in the number of stamens, each flower having often six, and sometimes seven. K.

CHELIDONIUM majus. Celandine. 11. * AQUILEGIA vulgaris. Columbine. RANUNCULUS bulbosus. Bulbous Crowfoot. AJUGA reptans. Common Bugle. GERANIUM robertianum. Stinking Crane's-bill. (12) GENISTA anglica. Needle Green-weed. . . TRIFOLIUM procumbens. Hop Trefoil. a LOTUS corniculatus. Common Bird's-foot Trefoil.

CAREX pulicaris. Flea Carex. ----- arenaria. Sea Carex.

QUERCUS Robur.
QUERCUS pedunculata. Willd. Common British Oak.

VALERIANA rubra. Red Valerian. 77 12. SCIRPUS caspitosus. Scaly-stalked Club-rush. a MYOSOTIS scorpioides. Mouse-ear Scorpion-grass. POLEMONIUM caruleum. Jacob's Ladder.

* CONVALLARIA multiflora. Common Solomon's Seal RUMEX Acetosella. Sheep's Sorrel.

* RUBUS arcticus. Dwarf crimson Bramble.

RANUNCULUS Flammula. Lesser Spear-wort. THLASPI perfoliatum. Perfoliate Shepherd's Purse.

COCHLEARIA Armoracia. Horse-radish.

CORONOPUS Ruellii.
COCHLEARIA Coronopus. Linn. Common Wart-cress

ERYSIMUM Barbarea. Yellow Rocket.

FUMARIA claviculata. White climbing Fumitory.

CAREX vulpina. Great spiked Carex.

URTICA dioica. Great Nettle.

13. SAGINA procumbens. Procumbent Pearl-wort. Rock 3/ ALLIUM ursinum. Broad-leaved Garlick. * CONVALLARIA majalis. Lily of the Valley. SCLERANTHUS annuus. Annual Knawel. RANUNCULUS repens. Creeping Crowfoot. ORNITHOPUS perpusillus. Common Bird's-foot.

TRIFOLIUM pratense, Common purple Trefoil.

⁽¹²⁾ Rarely varies to white.

SCIRPUS triqueter. Triangular Club-rush. ALOPECURUS pratensis. Meadow Fox-tail-grass. SAGINA erecta. Upright Pearl-wort. GALIUM cruciatum. Crosswort. Ite VALANTIA cruciata. Linn. LYSIMACHIA nemorum. Wood Loosestrife. HYDROCOTYLE vulgaris. Marsh Pennywort. * STAPHYLEA pinnata. Bladder-nut-tree. BERBERIS vulgaris. Barberry. (13) * VACCINIUM Vitis Idaa. Red Whortle-berry. STELLARIA uliginosa. Bog Stitchwort. ARENARIA peploides. Sea Sandwort. * PAPAVER cambricum. Yellow Poppy. ERYSIMUM officinale. Common Hedge-mustard. SPARTIUM scoparium. Common Broom. LATHYRUS Nissolia. Crimson Lathyrus. HIERACIUM Pilosella. Mouse-ear Hawkweed. URTICA weens. Small Nettle. ARUM maculatum. (Spadice purpureo.) Cuckowpint. STELLARIA graminea. Lesser Stitchwort. * DORONICUM Pardalianches. Great Leopard's-bane. CAREX acuta. Slender-spiked Carex. ----- riparia. Great common Carex. VERONICA Beccabunga. Brooklime. 16. & MYOSOTIS scorpioides. Mouse-ear Scorpion-grass. EUPHORBIA Peplus. Petty Spurge.

SISYMBRIUM Nasturtium. Water Cress.

SISYMBRIUM palustre. Willd. Annual water Rocket
HOLCUS avenaceus.
AVENA elatior. Linn. Oat-like Soft-grass.
POA trivialis. Roughish Meadow-grass.
GALIUM saxatile. Smooth heath Bed-straw.

⁽¹³⁾ It would be worth while to ascertain by experiment, whether the very probable conjecture of Dr. Smith respecting the fructification of this plant have any foundation in fact. K.

16. BUNIUM Bulbocastanum. Great Earth-nut.

HOTTONIA palustris. Water Violet.

* CONVALLARIA Polygonatum. Angular Solomon's Seal.

VICIA sativa. Common Vetch.

TRIFOLIUM minus. Lesser yellow Trefoil.

17. CORNUS sanguinea. Wild Cornel-tree.

ILEX Aquifolium. Holly-tree.

* LONICERA Xylosteum. Upright Honeysuckle.

VACCINIUM Oxycoccus. Cranberry.

POTENTILLA reptans. Common creeping Cinquefoil.

GEUM urbanum. Common Avens.

RANUNCULUS arvensis. Corn Crowfoot.

CAREX ovalis. Oval-spiked Carex.

18. VERONICA spicata. Spiked Speedwell.

----- officinalis. Male Speedwell.

LOLIUM perenne. Ray-grass.

SCANDIX Anthriscus. Rough Chervil.

ÆGOPODIUM Podagraria. Gout-weed.

RUMEX obtusifolius. Broad-leaved Dock.

GERANIUM dissectum. Jagged-leaved Crane's-bill.

TRIFOLIUM repens. White Trefoil.

ANTHEMIS Cotula. Stinking May-weed.

- 19. * ALLIUM Schanoprasum. Chive Garlick.
 - * CISTUS Helianthemum. Common dwarf Cistus.
 - * ARISTOLOCHIA Clematitis. Common Birth-wort.
 - * RHODIOLA rosea. Rose-root.
- 20. PARIETARIA officinalis. Pellitory of the Wall. SYMPHYTUM officinale. Common Comfrey.

* RHAMNUS Frangula. Alder Buckthorn.

* PYRUS Aria.

CRATEGUS Aria. Linn. White Beam-tree.

CRAMBE maritima. Sea Kale.

GERANIUM sanguineum. Bloody Crane's-bill.

MALVA rotundifolia. Dwarf Mallow. her

- * JUNIPERUS communis. Common Juniper.
- 21. SCIRPUS palustris. Marsh-creeping Club-rush.

22. FESTUCA pratensis. Meadow Fescue-grass, JUNCUS effusus. Soft Rush.
LYCHNIS Flos-Cuculi. Meadow Lychnis.
THYMUS Serpyllum. Wild Thyme.
SENECIO aquaticus. Marsh Ragwort.
ORCHIS Morio. Meadow Orchis.

SCIRPUS maritimus. Salt-marsh Club-rush. 23. POA fluitans. Flote Meadow-grass. FESTUCA fluitans. Linn. POA rigida. Hard Meadow-grass. DACTYLIS glomerata. Rough Cock's-foot-grass. PLANTAGO Coronopus. Buck's-horn Plantain. POTAMOGETON natans. Broad-leaved Pond-weed. SANICULA europea. Wood Sanicle. JUNCUS maritimus. Lesser sharp Rush. B JUNCUS acutus. Linn. POLYGONUM Bistorta. Great Bistort. B LYCHNIS dioica. White Campion. (14) HEDYPNOIS hispida. LEONTODON hispidum. Linn. Rough Hedypnois. APARGIA hispida. Willd. TAMUS communis. Black Bryony.

⁽¹⁴⁾ LYCHNIS vespertina of Sibthorp and Symons, so called from ital

- 24. POTAMOGETON gramineum. Grassy Pond-weed. (ENANTHE crocata. Hemlock Water Dropwort. ALLIUM vineale. Crow Garlick. ORCHIS maculata. Spotted palmate Orchis.
- 25. AIRA aquatica. Water Hair-grass.
 - * BRIZA maxima. Great Quake-grass. (Hull.)
 - * ALCHEMILLA alpina. Alpine Ladies-mantle.
 - * CYNOGLOSSUM officinale. Common Hounds-tongue. HERACLEUM Sphondylium. Common Cow-parsnep.
 - * SAXIFRAGA palmata. Palmate Saxifrage.

 SCROPHULARIA nodosa. Knotty-rooted Fig-wort. RAPHANUS Raphanistrum. Wild Radish. Wild Radish. TRAGOPOGON porrifolius. Purple Goat's-beard.

 OPHRYS ovata. EPIPACTIS ovata. Swartz.
- 26. SCIRPUS setaceus. Least Club-rush.

 SANGUISORBA oficinalis. Great Burnet.

 LITHOSPERMUM arvense. Corn Gromwell.

 VIBURNUM Opulus. Common Guelder-rose.

 PEPLIS Portula. Water Purslane.

 TRIGLOCHIN palustre. Marsh Arrow-grass.

 FUMARIA capreolata. Ramping Fumitory. (15)

 HIERACIUM murorum. Wall Hawkweed.
- 27. AVENA fatua. Wild Oat.
 - * RHAMNUS catharticus. Purging Buckthorn. / HYDROCOTYLE inundata. SISON inundatum. Linn. } Floating White-rot.
 - * MESPILUS germanica. Common Medlar.
 - * SPIRÆA Filipendula. Common Dropwort.
 - * GERANIUM columbinum. Long-stalked Crane's-bill.
- 28. AIRA præcox. Early Hair-grass.
 - * ASPARAGUS officinalis. Common Asparagus.
 - * SAXIFRAGA granulata. White Saxifrage.

⁽¹⁵⁾ Very common in the hedges in many places in this neighbourhood.

28. * CISTUS marifolius. Hoary dwarf Cistus. RANUNCULUS parvulus. Little upright Crowfoot. * ERODIUM moschatum. Musky Stork's-bill. GERANIUM moschatum. Linn. * VICIA lathyroides. Spring Vetch. - " t . - - !

29. VERONICA Anagallis. Water Speedwell. CYNOSURUS cristatus. Crested Dog's-tail-grass. GALIUM uliginosum. Rough marsh Bed-straw. SOLANUM Dulcamara. Woody Nightshade. (16) CHÆROPHYLLUM temulentum. Rough Cow-parsiey. COTYLEDON Umbilicus. B COTYLEDON Umbilicus. Linn. Common Navel-wort. COMARUM palustre. Marsh Cinquefoil. PAPAVER Rhaus. Common red Poppy. Pat. DIGITALIS purpurea. Purple Fox-glove. a'ORCIIIS bifolia. Butterfly Orchis.

AVENA pratensis. Narrow-leaved Oat-grass. ANAGALLIS arvensis. Scarlet Pimpernel. * EUONYMUS europaus. Common Spindle-tree. DAUCUS Carota. Wild Carrot. ACORUS Calamus. Sweet Flag. * TOFIELDIA palustris. ANTHERICUM calyculatum. Linn. Scottish Asphodel.

HELONIAS borealis. Willd. RESEDA Lateola. Dyer's-weed. NYMPHÆA lutea. Yellow Water-lily.

* HESPERIS inodora. Scentless Dame's-violet.

OROBUS tuberosus. Common Orobus.

ALOPECURUS bulbosus. Bulbous Fox-tail-grass. MELICA uniflora. Wood Melic-grass. * POA bulbosa. Bulbous Meadow-grass.

⁽¹⁶⁾ That the berries of this plant possess highly poisonous qualities is well known. That the stems are not entirely destitute of narcotic properties is equally true. In one instance where a very large dose had been swallowed, paralysiz of the tongue and fauces, with diminished sensibility of the retina was the consequence. Its effects were not entirely removed for several days. K.

31. * PINGUICULA vulgaris. Common Butterwort.

PLANTAGO maritima. Sea Plantain.

POTAMOGETON densum. Close-leaved Pond-weed.

* LITHOSPERMUM officinalc. Common Gromwell.

LYCOPSIS arvensis. Small Bugloss.

MEUM athamanticum.

ATHAMANTA Meum. Linn. ETHUSA Meum. Willd.

JUNCUS acutus. Great sharp Rush.

---- conglomeratus. Common Rush.

RUMEX sanguineus. Bloody-veined Dock.

SILENE maritima.

β CUCUBALUS Behen. Linn. Sea Campion.

ARENARIA marina.

\$\beta\$ ARENARIA rubra. Lir

Sea spurrey Sandwort.

RUBUS idaus. Raspberry.

PAPAVER Argemone. Long rough headed Poppy.

----- dubium. Long smooth headed Poppy.

NYMPHÆA alba. White Water-lily.

CARDUUS lanceolatus. Spear Thistle.

CHRYSANTHEMUM Leucanthemum. Great white Ox-eye.

ORCHIS latifolia. Marsh Orchis.

ARUM maculatum. (spadice albo) Cuckow-pint. (17)

SIXITH MONTH (June).

 SONCHUS arvensis. Corn Sow-thistle. ACHILLEA Millefolium. Common Yarrow.

2. RUMEX acutus. Sharp Dock.

POLYGONUM aviculare. Knotgrass.

* ALYSSUM sativum.
MYAGRUM sativum. Linu. } Gold of Pleasure.

3. BRIZA media. Common Quaking-grass.

⁽¹⁷⁾ I have observed that the inflorescence of those plants in which the Spadix is white, uniformly takes place later than when it is purple.

- 3. RUMEX crispus. Curled Dock.

 LATHYRUS latifolius. Broad-leaved Lathyrus.
- 4. CAUCALIS nodosa.
 TORDYLIUM nodosum. Linn.

 * SEDUM rupestre. Rock Stonecrop.
 RUBUS cassius. Dewberry.
 - ----- corylifolius. Hasel-leaved Bramble.
 TRAGOPOGON pratensis. Yellow Goat's-beard.
 ANTHEMIS arvensis. Corn Chamonnile.
 * CALENDULA arvensis. Marigold. (Hull.) (18)
 CAREX distans. Loose Carex.
- 5. SPERGULA nodosa. Knotted Spurrey.
 SISYMBRIUM amphibium. Great water Rocket.
 LEMNA minor. Lesser Duck-weed.
- 6. * SALVIA verbenaca. Wild English Clary. SAMBUCUS nigra. Dwarf Elder.

 MALVA sylvestris. Common Mallow. St. A. FAGUS Castanea. Chesnut-tree.
- 7. TOLIUM perenne. Ray-grass.
 TRITICUM repens. Couch-grass.
 * SAXIFRAGA stellaris. Hairy Saxifrage.
 * SILENE acaulis. Moss Campion.
- 8. PLANTAGO media. Hoary Plantain.

 JUNCUS uliginosus. Little bulbous Rush.

 * ANTIRRHINUM majus. Great Snapdragon.
- 9. * VERONICA saxatilis. Blue rock Speedwell. *alpina. Alpine Speedwell.

SCHOENUS albus. White-headed Bog-rush.

- * POA alpina. Alpine Meadow-grass.
- * STIPA pennata. Downy Feather-grass.
- * CAMPANULA glomerata. Clustered Bell-flower.

⁽¹⁸⁾ It is not uncommon for the flowers of this plant to become double; its multiplication arising from a conversion of the tubular into ligulate florets. It is worthy of remark that its fertility is increased in proportion to its multiplication. It is apt to become proliferous. K.

- * ATROPA Belladonna. Deadly Nightshade. 9. LONICERA Periclyminum. Common Honeysuckle,
 - * ANGELICA Archangelica. Garden Angelica.
 - * IMPERATORIA Ostruthium. Masterwort.

RUMEX aquaticus. Great water Dock.

ALISMA Plantago. Greater water Plantain. "

ERICA tetralix.

ERICA botuliformis. Salisbury. Cross-leaved Heath. (19)

- * SEDUM anglicum. English Stoneerop.
- * OXALIS corniculata. Yellow procumbent Wood-Sorrel. ROSA arvensis. White Dog-rose.

RUBUS fruticosus. Common Bramble.

- 10. FESTUCA elatior. Tall Fescue-grass. ELYMUS europæus. Wood Lyme-grass. EPILOBIUM palustre. Round-stalked marsh Willow-herb. GENISTA tinctoria. Dyer's Green-weed. SONCHUS oleraceus. Common Sow-thistle. CREPIS tectorum. Smooth Hawk's-beard.
- * LAGURUS ovatus. Hare's-tail-grass. 11. ÆTHUSA Cynapium. Fool's-Parsley. * EPILOBIUM angustifolium. Rose bay Willow-herb. MELAMPYRUM pratense. Common yellow Cow-wheat. HIERACIUM paludosum. Succory-leaved mountain Hawk-weed.
- VALERIANA officinalis. Great wild Valerian. LOLIUM arvense. White Darnel. HYOSCYAMUS niger. Common Henbane. CONIUM maculatum. Common Hemlock.
 - * ORNITHOGALUM pyrenaicum. Spiked Star of Bethlehem.
 - * DIANTHUS deltoides. Maiden Pink.
 - ----- cæsius. B DIANTHUS virgineus. Linn. Mountain Pink.

ARENARIA serpyllifolia. Thyme-leaved Sandwort.

Starting to a surprise and the or

⁽¹⁹⁾ All the species of Erica (E. Dabeoci excepted, and this I presume, will scarcely be considered as belonging to the genus) not unfrequently vary to white. 1 - Alangeli of the niverse is the site

- 12. SPIRÆA Ulmaria. Meadow-sweet.

 LINNÆA borealis. Two-flowered Linnæa.

 MEDICAGO polymorpha. Heart Medic.

 CARDUUS pratensis. Meadow Thistle.
- 14. AGROSTIS Spica-venti. Silky Bent-grass.

 FESTUCA ovina. Sheep's Fescue-grass.

 BROMUS erectus. Upright perennial Brome-grass.

 POTAMOGETON lucens. Shining Pond-weed.

 RUPPIA maritima. Sea Ruppia.

 ROSA spinosissima. Burnet Rose.

 OROBANCHE major. Greater Broom-rape.

 MEDICAGO sativa. Lucerne.
- 15. * POA flexuosa. Zigzag Meadow-grass.
 - * ----- casia. Sea-green Meadow-grass.
 - *---- nemoralis. Wood Meadow-grass.
 EPILOBIUM tetragonum. Square-stalked Willow-herb.
 BUTOMUS umbellatus. Flowering Rush.
 - * LYCHNIS Viscaria. Red German Catchfly.
 - * HEDYSARUM Onobrychis. Saintfoin. 4-4
 - * IHERACIUM auranticum. Orange Hawkweed.
- 16. CIRCEA lutetiana. Common Enchanter's Nightshade.

⁽²⁰⁾ Sometimes, although rarely, varies to purple.

- Common Cat's-tail-grass. PHLEUM pratense. 16.
 - Creeping Fescue-grass. * FESTUCA rubra.
 - Spiked Fescue-grass. * ----- loliacea.

Smooth rye Brome-grass. BROMUS seculinus.

Sheep's Scabious. JASIONE montanu.

* APIUM graveolens. Wild Celery.

ROSA caninu. Common Dog-rose. Programme 11X

* POTENTILLA finticosa. Shrubby Cinquefoil.

MENTHA arvensis. Corn Mint.

PRUNELLA vulgavis. Self-heal.

HYPERICUM quadrangulum. Square St. John's Wort.

SENECIO Jacobea. Common Ragwort.

BRYONIA dioica. Red-berried Bryouv.

AIRA cristata. Crested Hair-grass. 17.

POTAMOGETON perfoliatum. Perfoliate Pond-weed.

VICIA sepium. Bush Vetch.

GNAPHALIUM gullicum. Narrow-leaved Cud-weed. FILAGO gallica. Linn.

SENECIO sylvaticus. Mountain Groundsel.

BROMUS mollis. Soft Brome-grass. 13.

GALIUM Mollugo. Great hedge Bed-straw.

JUNCUS articulatus. Jointed Rush.

ALISMA ranunculoides. Small water Plantain.

SEDUM acre. Biting Stonecrop.

RESEDA lutea. Wild Mignonette.

EUPHORBIA helioscopia. Sun Spurge.

CREPIS biennis. Rough Hawk's-beard.

HYOSERIS minima. Swine's Succory.

ARUNDO Calamagrostis. Small Reed.

ROSA rubiginosa. Sweetbriar.

* PAPAVER somniferum. White Poppy.

SCUTELLARIA galericulata. Common Skull-cap.

* GERANIUM prateuse (flore albo). Crowfoot-leaved Crane's-bill.

TRIFOLIUM officinale. TRIFOLIUM Melilotus officinalis. Liun.

----- striatum. Soft knotted Trefoil.

19. * GNAPHALIUM luteo-album. Jersey Cud-weed.

20. LIGUSTRUM vulgare. Privet.

* ARUNDO colorata.
PHALARIS arundinacea. Linn. } Canary Reed-grass.

GALIUM palustre. White-water Bed-straw.

LINUM usitatissimum. Common Flax.

----- catharticum. Purging Flax.

* SILENE Armeria. Lobel's Catch-fly.

EUPHORBIA exigua. Dwarf Spurge.

† ----- paralia. Sea Spurge.

THALICTRUM flavum. Common Meadow-rue.

RANUNCULUS Lingua. Great Spear-wort.

SCROPHULARIA aquatica. Water Fig-wort.

ERVUM hirsutum. Hairy Tare.

HYPERICUM androsamum. Tutsan St. John's Wort.

MATRICARIA Chamomilla. Wild Chamomile.

TYPHA angustifolia. Lesser Reed-mace.

HYDROCHARIS Morsus rana. Common Frog-bit.

21. *VERONICA fraticulosa. Flesh-coloured shrubby Speed-

SCIRPUS Holoschoenus. Round Cluster-headed Club-rush.

FESTUCA duriuscula. Hard Fescue-grass.

LOLIUM temulentum. Bearded Darnel.

LYSIMACHIA Nummularia. Creeping Loosestrife.

CAMPANULA rotundifolia. Round-leaved Bell-flower.

HERNIARIA glabra. Smooth Rupture-wort.

GERANIUM pratense. Crowfoot-leaved Crane's-bill.

HYPERICUM humifusum. Trailing St. John's-Wort.

22. * CIRCÆA alpina. Mountain Enchanter's Nightshade. + ARUNDO arenaria. Sea Reed.

SEDUM album. White Stonecrop.

23. CHENOPODIUM urbicum. Upright Goosefoot.
POLYGONUM Convolvolus. Climbing Buckwheat.
HYPERICUM pulchrum. Upright St. John's-Wort.
LAPSANA communis. Nipple-wort.
PYRETHRUM Parthenium.
MATRICARIA Parthenium. Linn. Common Feverfew.

- 24. VERONICA scutellata. Narrow-leaved Marsh Speedwell GALIUM verum. Yellow Bed-straw.
 - ANTIRRHINUM Linaria. Common yellow Toadflax.
 - * HYPERICUM hirsutum. Hairy St. John's Wort.
- 25 HORDEUM maritimum. Sea Barley.
 - PLANTAGO major. Greater Plantain.
 - * IMPATIENS Noli me tangere. Yellow Balsam.
 - SOLANUM nigrum. Common Nightshade. (21)
 - * ERYNGIUM campestre. Field Eryngo.
 - SIUM latifolium. Broad-leaved Water Parsnep.
 - AGROSTEMMA Githago. Corn Cockle. 1 /
 - * DELPHINIUM Consolida. Field Larkspur.
 - * ADONIS autumnalis. Corn Adonis.
 - † BLOTUS corniculatus. Common Bird's-foot 'Trefoil
 - CHRYSANTHEMUM segetum. Yellow Ox-eye.
 - ATRIPLEX pedunculata. Pedunculated sea Orache.
- 26. HORDEUM pratense. β HORDEUM murinum. Linn. } Meadow Barley.

 CICUTA virosa. Water Hemlock.

 'TRIFOLIUM ochroleucum. Sulphur coloured Trefoil.

 α CENTAUREA nigra. Lesser Knapweed.
- 27. BORAGO officinalis. Common Borage.
 - DATURA Stramonium. Thorn Apple.

ŒNANTHE fistulosà. Common Water Dropwort.

CHLORA perfoliata.
GENTIANA perfoliata. Linn. } Yellow Centaury. (22)

POLYGONUM Fagopyrum. Buckwheat.

MARRUBIUM vulgare. White Horehound.

HYPERICUM perforatum. Perforated St. John's Wort.

⁽²¹⁾ Too much pains cannot be taken to extirpate this poisonous weed, the berries of which have an appearance likely to tempt the appetites of children. The situations in which it is most frequently found near Warrington, are the gardens at the western end of the town and the road side at Lower Walton. K.

⁽²²⁾ This plant may be added to the list of those which are usually denominated equinoctial; its expansion commencing at 7 o'clock A. M. and ending at 2 o'clock P. M. K.

HORDEUM murinum. Wall Barley. SAMBUCUS Ebulus. Dwarf Elder. NAR'THECIUM ossifragum. ANTHERICUM ossifragum. Linn. Lancashire Asphodel VERBENA officinalis. Common Vervain. CARDUUS arvensis. SERRATULA arvensis. Linn. { Creeping Thistle. EUPATORIUM cannabinum. Hemp-agrimony. TYPHA latifolia. Broad-leaved Reed-mace. POLYGONUM Persicaria. Spotted Persicaria. 29. SEMPERVIVUM tectorum. Common Houseleek. * ACTÆA spicata. Herb Christopher. TILIA europæa. Common Lime-tree. GALEOPSIS versicolor. Large-flowered

GALEOPSIS cannabina. Willd.) + SISYMBRIUM Sophia. Flix-weed.

B GALEOPSIS Tetrahit. Linn.

30. CHENOPODIUM nubrum. Red Goose-foot.

album. White Goose-foot.

BSAMBUCUS nigra. Common Elder.

† PYROLA minor. Lesser Wintergreen. (23).

SEDUM reflexum. Yellow Stonecrop.

* POTENTILLA alba. White Cinquefoil.

* SIBTHORPIA europæa. Cornish Moneywort.

GNAPHALIUM germanicum. FILAGO germanica. Linn. Common Cudweed.

B CENTAUREA nigra. Lesser Knapweed.

SPARGANIUM simplex. Unbranched upright Bur-reed.

Hemp-nettle.

SEVENTH MONTH (July).

1. VERBASCUM Thapsus. Great Mullein. Ma

* LIGUSTICUM scoticum. Scottish Lovage.

STACHYS arvensis. Corn Woundwort.

† 7 ONONIS arvensis.

ONONIS repens. Linn.

Rest-harrow.

⁽²³⁾ This rare plant was brought to me from a rock near the waterfall at Ambleside, Westmoreland.

- VALERIANA dentata. Oval-fruited Corn Sallad. + ERYNGIUM maritimum. Sea Eryngo. PHELLANDRIUM aquaticum. Water Hemlock. Yellow Horned-† GLAUCIUM luteum. CHELIDONIUM Glaucium. Linn. † ANTHYLLIS vulneraria. Kidney Vetch. † VICIA sylvatica. Wood Vetch.
- * ----- Cracca. Tufted Vetch-FESTUCA gigantea. Tallest Fescue-grass. BROMUS giganteus. Linn.

CONVOLVOLUS sepium. Great Bindweed. AGRIMONIA eupatoria. Common Agrimony. STACHYS palustris. Marsh Woundwort.

a ONONIS arvensis. Rest-harrow.

POA maritima. Sea Meadow-grass. ANAGALLIS tenella. LYSIMACHIA tenella. Linn. Bog Pimpernel. GENTIANA Centaurium. Linn. Common Centaury.

LYSIMACHIA vulgaris. Yellow Loosestrife. 5. + GERANIUM sylvaticum. Wood Crane's-bill. MALVA moschuta. Musk Mallow. + HYPERICUM montanum. Mountain St. John's Wort.

+ BCIRCÆA ulpina. Mountain Enchanter's CIRCÆA intermedia. Ehrhart. \ Nightshade. CONVOLVOLUS arrensis. Small Bindweed.

SIUM nodiflorum. Procumbent Water Parsnep.

PIMPINELLA saxifraga. Common Burnet-Saxifrage.

* OENOTHERA biennis. Biennial Oenothera. (E. Bor.)

BALLOTA nigra. Stinking Black Horehound.

LEONURUS Cardiaca. Motherwort.

* HYPERICUM calycinum. Great-flowered St. John's Wort. (E. Bor.)

BIDENS tripartita. Trifid Bur Marygold. CENTAUREA Scabiosa. Greater Knapweed.

SCABIOSA arrensis. Field Scabious. CAMPANULA latifolia. Giant Bell-flower. 7. EPILOBIUM roseum. Pale Smooth-leaved Willow-herb.
TEUCRIUM Scorodonia. Wood Germander.

\$\beta\$ ONOMIS arvenis.
\$\beta\$ ONOMIS spinosa. Linn. } Rest-harrow.

CARLINA vulgaris. Common Carline.

GNAPHALHIM waygavitagetty. Appriser Codesale.

* GNAPHALIUM margaritaceum. American Cudweed. ACHILLEA Ptarmica. Sneeze-wort.

* CAMPANULA Trachelium. Nettle-leaved Bell-flower.

* CHENOPODIUM olidum.
CHENOPODIUM Vulvaria. Linn. Stinking Goose-foot.

* SWERTIA perennis. Marsh Felwort.

SELINUM palustre. Marsh Milky Parsley.

CAUCALIS Anthriscus.
TORDYLIUM Anthriscus. Linn. Upright Hedge Parsley

JUNCUS bufonius. Toad Rush.

EPILOBIUM hirsutum. Great hairy Willow-herb.

ARENARIA tenuifolia. Fine-leaved Sand-wort.

9. MELICA carulea.
AIRA carulea. Linn. Purple Melic-grass.
GALEOPSIS Tetrahit. Common Hemp-nettle.
ERVUM tetraspermum. Smooth Tare.
CENTAUREA Calcitrapa. Common Star-thistle.
† ORCHIS conopsea. Aromatic Orchis.

* CARDUUS marianus, Milk Thistle.

10. SAMOLUS Valerandi. Brookweed.

† CONVOLVOLUS Soldanella. Sea Bindweed.

† SALSOLA Kali. Prickly Saltwort.

LYTHRUM Salicaria. Purple Loosestrife. (24)

BETONICA officinalis. Wood Betony.

† BUNIAS Cakile.

CAKILE maritima. Willd. Sea Rocket.

CICHORIUM Intybus. Wild Succory.

⁽²⁴⁾ A singular variety of this is sometimes found, which differs in the foliation, its leaves being alternate, ovate, and acuminate. This is not accidental but permanent. K.

- SERAPIAS latifolia. SERAPIAS Helleborine. Linn. > Broad-leaved Helleborine EPIPACTIS latifolia. Swartz.) SERAPIAS palustris. y SERAPIAS longifolia. Linn. > Marsh Helleborine. EPIPACTIS palustris. Swartz. † ATRIPLEX laciniata. Frosted sea Orache.
- Manured Canary-grass. * PHALARIS canariensis. 11. EPILOBIUM parviflorum. B EPILOBIUM hirsutum. Linn. Willow-herb. * DIANTHUS Caryophyllus. Clove Pink.
 - * SERRATULA alpina. Alpine Saw-wort.
 - 12. * IRIS fatidissima. Stinking Iris. SCABIOSA succisa. Devil's-bit Scabious. JUNCUS bulbosus. Round-fruited Rush. * DIAN'THUS prolifer. Proliferous Pink. STACHYS sylvatica. Hedge Woundwort. ASTER Tripolium. Sea Star-wort.

PYRETHRUM maritimum. Sea Feverfew. MATRICARIA maritimum. Linn.

- 13. LYCOPUS europæus. Water Horehound. a DIPSACUS fullonum. Linn. Wild Teasel. DIPSACUS sylvestris. * CAMPANULA Rapunculus. Rampion Bell-flower. VERBASCUM Blattaria. Moth Mullein. SIUM anguistifolium. Narrow-leaved Water Parsnep. PIMPINELLA magna. Great Burnet-Saxifrage. * STRATIOTES aloides. Water Soldier. GNAPHALIUM minimum. Least Cudweed.
- * LYSIMACHIA thyrsiflora. Tufted Loosestrife. ANETHUM Faniculum. Common Fennel. STATICE Limonium. Sea Layender. DROSERA rotundifolia. Round-leaved Smi-dew. (25) --- longifolia. Long-leaved Sun-dew.

14.

⁽²⁵⁾ The following accurate description of the Phanomenon exhibited by this Plant, may be considered worthy of insertion in this place.

[&]quot; Ses seuilles sont bordées de poils terminés par une petite goutte de liqueur. Quand un insecte se pose sur le disque d'une feuille, elle se ferme comme x lee a e la trela ? 17

14. † β GALEOPSIS Ladanum. Red Hemp-nettle.

* LAVATERA arborea. Sea Tree-mallow.

HYPERICUM elodes. Marsh St. John's Wort.

HEDYPNOIS autumnalis.

LEONTODON autumnalis. Linn.

APARGIA autumnalis. Willd.

* INULA Helenium. Elecampane.

15. SAXIFRAGA Hirculus. Yellow Marsh Saxifrage. (26)
BARTSIA viscosa. Yellow viscid Bartsia. (27)
ARTEMISIA vulgaris. Mugwort.
+ CONYZA squarrosa. Plowman's Spilzenard.

16. POA procumbens. Procumbent sea Meadow-grass,
† DIPSACUS pilosus. Small Teasel.
* NEPETA Cataria. Cat-mint,
SONCHUS palustris. Tall marsh Sow-thistle.
TANACETUM vulgare. Common Tansy.

une bourse à jetons dont on tire les cordons, et l'insecte meurt, tout couvert du suc visqueux qui s'echappe do l'extremité des poils. C'est un spectacle aussi bizarre qu'il est etonnant, de voir, comme j'ai eu plusieurs fois occasion de l'observer, presque toutes les feuilles de cette petite plante formant des prisons enduites de glu, où sont retenus quelques insectes, tels que des fourmis ou des petites mouches. Ainsi la rencontre d'un gazon de rossolis est aussi terrible, pour ces légions de créatures fobles et misèrables, que l'explosion d'un feu souterrain pour les peuplades humaines; mais ces desordres ont leur limites; et, tandis que les individus sont en proie à mille dangers, les espéces ne cessent pas de se propager, et ces innombrables races d'êtres ephémeres no sont pas meins que la race humaine sous la main protectrice de la Nature."

MIRBEL Traite d'Anatomie

ct de Physiologie Vegetales

Tom. 1. p. 265.

- (26) Observed upon Knutsford moor; the habitat already pointed out.
- (27) This rare plant has been found at Ormskirk, South-port, Allerton, Speke, Hale, and Ditton in Lancashire, and on Bartington heath in Cheshire, by different members of the Estanical Society of Warrington.

- 17. † PARNASSIA palustris. Grass of Parnassus. HUMULUS Lupulus. Common Hop.
- 18. SEDUM Telephium. Orpine.

 * SPIRÆA salicifolia. Willow-leaved Spiræa.

 GALEOPSIS villosa. Downy Hemp-nettle.

 α ONONIS arvensis (flore albo). Rest-harrow. (28)

 INULA dysenterica. Common Flea-bane. (29)

 ATRIPLEX patula. Spreading halberd-leaved Orache.
 - 19. GENTIANA Pneumonanthe. Marsh Gentian. (30)
 ERICA vulgaris.
 CALLUNA vulgaris. Salisbury.

 BARTSIA Odontites.
 EUPHRASIA Odontites. Linn.
 PICRIS hieracioides. Hawkweed Ox-tongue.
 LACTUCA Scariola. Prickly Lettuce.
 ARTEMISIA maritima. Sea Wormwood.
 - 20. BROMUS sylvaticas. Slender wood Brome-grass,

 * HERNIARIA hirsuta. Hairy Rupture-wort.

 * ERICA vagans.
 ERICA vaga. Salisbury.

 POLYGONUM lapathifolium. Pale-flowered Persicaria.

 ANTHEMIS maritima. Sea Chamomile.
 - 21. * VERBASCUM Lychnitis. White Mullein. MENTHA sylvestris. Horse Mint.

⁽²⁸⁾ This variety, differing from α Ononis arrensis, not only in the whiteness of its petals, but in the colour of the stalk which is of a pale green instead of purple, is found near the pier, at Runcorn. K.

⁽²⁹⁾ This plant is said to have been occasionally useful in cases of Dysentery. As this disease, after an interval of more than 100 years, has of late appeared among us, it may perhaps, as a cheap remedy, be worthy of trial. It has an unpleasant smell, compared by some to that of soap; and a bitterish subacid taste. Its acrimony a little resembles that of Arum maculatum. K.

⁽³⁰⁾ A white-flowering variety of this plant, is said to have been found upon Bartington heath.

- 21. SOLIDAGO Virgaurea. Common Golden-rod. (31)
- 22. * CORIANDRUM sativum. Common Coriander.

 * TEUCRIUM Chamædrys. Common Germander.

 CINERARIA palustris. Marsh Flea-wort.
- 23. POLYGONUM Hydropiper. Biting Persicaria. MERCURIALIS annua. Annual Mercury.
- * SALVIA pratensis. Meadow Clary.
 * DIPSACUS fullonum. Manured Teasel.
 DROSERA anglica. Great Sun-dew.
 ANDROMEDA polifolia. Marsh Rosemary. (Second inflorescence.) (32)
 ARCTIUM Lappa. Burdock.
- 25. SENECIO viscosus. Stinking Groundsel.
- 26. SALICORNIA herbacea. Marsh Samphire. ERIGERON acre. Blue Flea-bane.
- 27. CAUCALIS infesta.
 SCANDIX infesta. Linn. Spreading hedge Parsley.
 SENECIO tenuifolius. Hoary Ragwort.
- 28. ANGELICA sylvestris. Wild Angelica.
 ANTHEMIS nobilis. Common Chamomile.
- 29. CHIRONIA Centaurium. (flore albo) Common Centaury. (33)

 MENTHA hirsuta.: Hairy Mint.
 LATHYRUS palustris. Marsh Lathyrus.
- 30. MENTHA rotundifolia. Round-leaved Mint. HYPOCHÆRIS radicata. Long-rooted Cat's-ear.
- 31. GENTIANA Amarella. Autumnal Gentian. STACHYS germanica. Downy Woundwort.

⁽³¹⁾ The powdered Root, in doses of half a dram, three times a day, has been found a powerful remedy in *Dropsy*. It has long been in use in cases of *Nephritis*, and with good effect. K.

⁽³²⁾ Repeated observation confirms the opinion that this species has two distinct periods of inflorescence in each year.

⁽³³⁾ This variety has been found in several parts of this neighbourhood. K.

31. HIERACIUM sabaudum. Shrubby broad-leaved Hawk-weed.

URTICA pilulifera. Roman Nettle.

EIGHTH MONTH (August).

streat. ", mined the one

- 1. SCUTELLARIA minor. Lesser Skull-cap.
- 2. GNAPHALIUM uliginosum. Marsh Cudweed.
- 3. * MENTHA viridis. Spear Mint.
- 4. GNAPHALIUM rectum. Upright Cudweed.
- 5. BIDENS cernua. Nodding Bur Marygold.
- 6. POLYGONUM minus. Small creeping Persicaria.
- 7. CLINOPODIUM vulgare. Wild Basil.

 LEONTODON palustre.

 LEONTODON lividus. Willd.

 Marsh Dandelion.
- 8. ARUNDO Phragmites. Common Reed.
- 9. MENTHA piperita. Pepper Mint.
- 10. ----- rubra. Tall red Mint.
- 11. * ORIGANUM vulgare. Common Marjoram.
- 12. ARUNDO epijegos. Wood Reed.
- 13. MENTHA odorata. Bergamot Mint.
- 14. UTRICULARIA vulgaris. Hooded Milfoil.
- 16. MENTHA Pulegium. Penny Royal.
- 17. SAPONARIA officinalis. Soapwort.
- 19. CHIRONIA pulchella. Dwarf branched Centaury.
- 20. CHENOPODIUM hybridum. Maple leaved Goose-foot.

 * HIERACIUM umbellatum. Narrow-leaved Hawkweed.

 † SERRATULA tinctoria. Common Saw-wort.
- 24. OPHRYS spiralis. NEOTTIA spiralis. Swartz. } Spiral Ophrys.
- 25. 7 BIDENS cernua.
 BIDENS minima. Linn. } Nodding Bur Marygold.

- 27. CORNUS sanguinea. Wild Cornel-tree (second inflorescence). (34)
- 28. SCLERANTHUS perennis. Perennial Knawel.

NINTH MONTH (September).

- 8. * COLCHICUM autumnale. Meadow Saffron.
- 10. * ARBUTUS Unedo. Strawberry-tree.
- 18. * CROCUS sativus. Saffron Crocus.
- 24.

 γ CALLITRICHE aquatica.

 CALLITRICHE autumnalis. Linn.

 Water Star-wort.

TENTH MONTH (October).

13. HEDERA Helix. Common Ivy. 157

⁽³⁴⁾ The time here mentioned was evidently the second period of inflorescence, but whether it is permanently the case as in Andromeda I do not know.

INDEX GENERUM.

The Genera printed in Italics occur only as synonyms.

			Page.	? Page.
	Α		- 5	Angelica - 21, 32.
ACER		-	10.	Anthemis - 15, 20, 31, 32.
Achillea	•		19, 28.	Anthericum - 18, 26.
Acorus -		-	18.	Anthoxanthum - 11.
Actæa			26.	Anthyllis 27.
Adonis -		-		Antirrhinum - 11, 20, 25.
Adoxa	_	-	7.	<i>Apargia</i> - 16, 30.
Ægopodium			15.	
Æthusa	~		19, 21.	Apium 23.
Agrimonia		-	27.	
Agrostemina		-	25.	
Agrostis	-		16, 22.	
Aira -		17	, 23, 28.	
Ajuga	-	-	13.	Arenaria 12, 14, 19, 21, 28.
Alchemilla	-		7, 8, 17.	Aristolochia - 15.
Alisma	-		21, 23.	{ Artemisia - 30, 31.
Allium		13	, 15, 17.	} Arum 14, 19.
Alnus -	•	-	6	Arundo - 23, 24, 33.
Alopecurus	-	14	, 16, 18	S Asarum - 9.
Alsine	-	-	6	Asparagus - 17.
Alyssum	-	-	19	Asperula 9.
Anagallis		-	18, 27	· Aster 29.
Anchusa	•	-	10	· \ Athamanta - 19.
Andromeda		-	10, 32	. Atriplex - 25, 29, 31.
Anemone			7,8	
Anethum	-	•	29	

INDEX.

B Page.	P
Ballota - 27.	(())
Bartsia 30, 31.	Colobin
Bellis - 5	Comarun
Berberis - 14	Conium
Betonica - 28	2 (1 1)
Betula - 6 0	S Count 1
Bidens - 27, 33.	Comme
Borago	Carinada
Brassica - 9.17	The state of the s
Briza 17, 10	(Canana
Dronius - 22, 23, 27, 31	/ Courles
Bryonia - 23.	Cotylodan
Bunias - 29	(Charal
Bunium - 11 15	Ount.
Butomus 22.	Channel
Buxus 8.	1 112
C C	5 (7)
Cakile 28.	(Truspelle -
Calendula - 20.	Cuanin
Callitriche - 8, 31.	Cynosurus - 18.
Caltha 7.	Dagfult
Calluna - 31.	/ Than i
Campanula 20, 24, 27, 28, 29.	1 Datama
Cardamine - 0, 8, 10.	Daucus - 25.
Carduus - 19, 22, 26, 28.	Delphinium - 25.
Carex 10, 11, 13, 14, 15, 16, 20.	Dentaria - 8.
Carlina - 28.	Dianthus 21, 29.
Carpinus 3.	Digitalis - 18.
Caucalis - 20, 28, 32.	Dipsacus - 29, 30, 32.
Centaurea 12,25,26,27,28.	Doronicum - 14.
Cerastium - 7, 9, 10, 11.	Draba - 6.
Cherophyllum - 9, 18.	Drosera - 29, 32.
Chara - 7, 9	E
Cheiranthus - 7.	Echium 8.
Chelidonium - 13, 27.	Elymus - 21.
Chenopodium 11,21,26,28,33.	Epilobium > 21, 22, 28, 29.
Chironia - 27, 32, 33.	Epinedium - 9.
Chlora - 25.	Epipactis 17, 29.
Chrysanthemum' - 19, 25.	Erica - 10, 21, 22, 31.
Chrysoplenium - 7.	Erigeron - 32.
Cichorium 28.	Eriophorum - 9.
Cicuta 25.	Erodium - 10, 18.
Cineraria - 32.	Ervum - 24, 28.
Circæa - 22, 24, 27.	Eryngium - 25, 27.
and the second s	Erysimum - 9, 13, 14.
Clinopodium - 33.	Euonymus - 18.

INDEX.

	Page.	$Pa_{\mathbf{g}}$	e.
Eupatorium	- 26.		3.
Euphorbia	- 9, 14, 23, 24.	Hypericum 23, 24, 25, 27, 3	0.
Euphrasia	22, 31.		2.
25ap 11 to 11	F	I	
Fagus -	- 10, 20.	Iberis 10, 1	2.
	21, 22, 23, 24, 27.	Ilex 1	5.
Filago	23, 26.	Impatiens - 2	25.
Fragaria	6, 9.	Imperatoria - 2	21.
Fraxinus	12.	Inula 30, 3	31.
Fritillaria	8.	Iris 16, 2	29.
Fumaria -	8, 10, 12, 13, 17.	XIxia	9.
	G	}J	
Galanthus	5.		23.
Galeobdolon	- 12.	Juncus 7,12,16,19,20,23,28,5	29.
Galeopsis 12	2,22,26,28,30,31.	Juniperus -	15.
Galium 8	3, 14, 18, 23, 24, 25.	}_ L	
Genista	- 13, 21.		31.
Gentiana -	10,25,27,31,32.		21.
Geranium 10	0, 12, 13, 15, 17, 18,	Lamium 5,	7.
	3,24,27.	{ Lapsana	24.
Geum	11, 15.	Lathyrus - 14, 20, 22,	
Glaucium	27.)	30.
Glaux -	12.	•	20.
Glechoma -	- 7.	Leontodon - 6, 16, 30,	
Gnaphalium	12,23,24,26,28,	Leonurus - 12,	
Andrew .	29,33.		10.
** . 1	H	, ,	26.
Hedera .	34.	1	24.
Hedypnois	- 16,30.	(~ .	22.
Hedysarum Helleborus	- 22. - 5, 8.	1 - 1 1	24.
Helonias	- 18.		
Heracleum	- 17.		
Herniaria	- 24,31.	4	
Hesperis	- 18.		
Hieracium	14, 17, 21, 22, 33.	/ _ ~ .	19.
Hippuris	- 10.	3 4 7 1	29.
Holcus	- 14, 16.		
Holosteum	- 10.		28.
Hordeum	- 25, 26.	S M	
Hottonia	- 15.	. Malva - 15, 20.	27.
Humulus	- 31.	. ¿ Marrubium 🕒	25.
Hyascinthus	- 11.	.; Matricaria - 24.	29.
Hydrochari	24.	. Medicago	22.
Hydrocotyl		. Melampyrum -	21.
Hyoscyamu	ıs - 21	. [Melica 18,	28.

INDE,X.

	Page.	4 3	Page.
Montha - 2	3, 31, 32, 33.	Polygonum	16, 19, 22, 24, 25,
Menyanthes	- 12.		26, 31, 32, 33.
Menziesia	- 10.	Populus	- 6, 7, 8.
Mercurialis -	7, 32.	Potamogeton	16, 17, 19, 22, 23.
Mespilus -	11, 17.	Potentilla	12,-15, 23, 26.
Meum -	19.	Prenanthes	
Montia -	- 7.	Primula	5, 7, 8.
Myagrum -	- 19.	Prunella	23,
Myosotis -	12, 13, 14.	Prunus	- 8, 9, 12.
Myrica	8.	Pulmonaria	
N		Pyrethrum	
Narcissus -		Pyrola	-, - 26.
Narthecium	26.	Pyrus -	9, 11, 12, 15.
Neottia , -	33.	{	Q
Nepeta -	- 30.	Quercus	13.
Nymphæa -	18, 19.	} D	R
, 0		Ranunculus	
Enanthe	17, 25.	En. L.	
Oenothera -	- 27.	Raphanus	17.
	26, 27, 28, 31.	Reseda .	- 18, 23. - 15, 17.
Ophrys -	17, 33.	Rhamnus	- 10, 17.
Orchis 10, 16, 17,	18, 19, 22, 28.	Rhodiola	- 16. - 15.
Origanum-	- 33.	Ribes -	7, 8, 9.
	. 8, 10, 21.	Rosa -	
Ornithopus	- 13.		13, 19, 20, 21.
Orobanche -	- 22.		, 13, 15, 19, 20, 21.
Orobus -	- 18. - 6, 21.	Ruppia	22.
Oxalis - P		Ruscus	_ '8.
	11 19 10 93	, reasons	S
Auron	14, 18, 19, 23. - 15.	Sagina .	- 11, 13, 14.
Parietaria	12.	Salicornia	32.
Paris -	31.	Salix .	- 6, 9, 10, 11.
Parnassia		Salsola, -	- 28.
Pedicularis	17.	Salvia -	- 20, 32.
Peplis - Phalaris -	24, 29.		- 20, 26.
Phalandrium -	27.		- 28.
Phleum -	. 23.		17.
Picris -	31.	Sanicula	16.
Pimpinella	27, 29.		_ 33.
Pinpineua Pinguicula	19.	2 Savifraga	11, 12, 17, 20, 30.
Plantago - 9	16, 19, 20, 25.		27, 29,
Poa 6, 14, 16, 18	0.90.92, 27, 30.) Scanuix	11, 15, 32.
Polemonium	_ 13.	Cochamas	20.
Polygala -	11.	Scilla	- 6, 7, 10, 11.
i diy Sala		-	

INDEX.

n 2	Page.
Page.	Taxus - 7.
Scirpus 13, 14, 15, 16, 17, 24.	Teucrium - 28, 32.
Scleranthus - 13, 34.	Thalictrum - 22, 24.
Scrophularia - 8, 17, 24.	Thlaspi - 7, 8, 9, 13.
Scutellaria - 23, 33.	16
Sedum 20, 21, 23, 24, 26, 31.	Thymus 10. Tilia 26.
Selinum 23.	Tofieldia 18.
Sempervivum - 26.	1 Ollelula
Senecio - 5, 16, 23, 32, 33.	, xo, ayean
Serapias 29.	Tormentilla - 12. Tragonogon - 17, 20.
Serratula - 26, 29, 33.	
Sherardia - 12.	10 17
Sibthorpia - 26.) 111610011111
Silene - 19, 20, 22, 24.	(I riticum
Sinapis 10.) 1 romus -
Sison 17.	Tunpa
Sisymbrium - 14, 20, 26.	Luillus
Sium - 25, 27, 29.	CA OR
Solanum - 18, 25.) Lypna Tr
Solidago 32.	- 00
Sonchus - 19, 21, 30.	6 7
Sorbus 12.	70 14 00
Sparganium - 26.	00
Spartium 14.	Officulation
Spergula - 16, 20.	
Spiræa - 17, 22, 31.	3
Stachys - 26, 27, 29, 32.) valuntia
Staphylea 14.	
Statice - 12, 29	0.0
Stellaria - 6, 9, 14) Y CI UCIIA
Stipa 20	
Stratiotes - 29	1
Swertia 28	· (VIUULIIUIII
Dy Impary value	. Vicia - 15, 18. 23, 27.
T	Vinca 5, 8.
T. COLLEGE	. Viola - 5, 6, 7, 8, 11.
Tanacetum - 30	. \chi_\text{Viscum} \text{-} \text{-} \text{8.}

THE END.

J. Haddock, Printer, Horse-Market, Warrington.